

400-544

6/13/2014

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

JUN 13 2014

Denise Dzialo
Chemtura Corporation
199 Benson Rd
Middlebury, CT 06749

Subject: Supplemental label for legume uses of Rancona 3.8 FS
Submission Date: 3/27/2014 with a resubmission 6/13/2014
Product Name: Rancona 3.8 FS
EPA Registration No.: 400-544
EPA Decision No.: 489588

Dear Ms. Dzialo,

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act is acceptable under FIFRA 3(c)(5).

The agency acknowledges the creation of a supplemental label, which covers the use of Rancona 3.8 FS on legumes. This new use was approved 3/20/2014; therefore, this supplemental label may not be distributed or used after 3/20/2017. A copy of this supplemental label has been stamped "Accepted" and will be put in our records for this product.

If you have any questions, please contact Lindsay Roe by phone at (703) 347-0506 or via email at roe.lindsay@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Tony Kish".

Tony Kish
Product Manager (22)
Fungicide Branch
Registration Division (7504P)

SUPPLEMENTAL LABEL

The supplemental label expires on 3/20/2017 and must not be used or distributed after this date.

	<h2>Rancona[®] 3.8 FS</h2> <h3>SEED TREATMENT FUNGICIDE</h3> <p>EPA Reg. No. 400-544</p> <h4>For Use on Legume Vegetables (Succulent or Dried)</h4>								
<table border="0"> <tr> <td>Active Ingredients: (% by weight)</td> <td></td> </tr> <tr> <td>ipconazole (CAS No. 125225-28-7).....</td> <td>40.7%*</td> </tr> <tr> <td>Other Ingredients:</td> <td>59.3%</td> </tr> <tr> <td>Total:.....</td> <td>100.000%</td> </tr> </table> <p>*Contains 3.77 pounds of active ingredient per gallon. *Contains 452 grams of active ingredient per liter.</p>		Active Ingredients: (% by weight)		ipconazole (CAS No. 125225-28-7).....	40.7%*	Other Ingredients:	59.3%	Total:.....	100.000%
Active Ingredients: (% by weight)									
ipconazole (CAS No. 125225-28-7).....	40.7%*								
Other Ingredients:	59.3%								
Total:.....	100.000%								
<p>KEEP OUT OF REACH OF CHILDREN</p> <p>CAUTION</p>									

ACCEPTED
 JUN 13 2014
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 400-544

THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL ON THE CONTAINER

Follow all applicable directions, restrictions and precautions on this supplemental labeling and the main EPA registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the federal label for this product must be in the possession of the user at the time of the pesticide application.

Read the label affixed to the container for Rancona 3.8 FS before applying. Use of Rancona 3.8 FS according to this labeling is subject to the use precautions and limitations imposed on the label affixed to the container for Rancona 3.8 FS.

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. Care must be exercised in the handling of treated seed. Do not use augers used for handling treated seed to move seed for feed, food or oil processing. Do not re-use bags from treated seed to handle food or feed products.

ALWAYS SHAKE OR STIR PRODUCT WELL BEFORE USE

Manufactured for:
 Chemtura Corporation
 199 Benson Road
 Middlebury, CT 06749

015A/

NOTE: This product is undyed. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately colored with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

SEED TREATMENT EQUIPMENT

Rancona 3.8 FS may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease protection and optimum performance. Seed should be sound and well-cured before treatment. Refer to the label rates below. Product requires dilution prior to application. Contact your local Chemtura representative or supplier for specific recommendations.

SEED TREATMENT: APPLICATION RATES

LEGUME VEGETABLES (SUCCULENT OR DRIED):

Bean (Lupinus) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (Vigna) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (Pisum) (includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; soybean; soybean (immature seed); sword bean.

For protection against seed rot, damping off and seedling blight including Fusarium (seed and soil borne) and Rhizoctonia solani. Also protects against general seed rots caused by saprophytic organisms such as Penicillium and Aspergillus, and protects against seed-borne Botrytis and Sclerotinia. Provides suppression of seed borne Anthracnose of dry bean. Apply 0.085 fl. oz. of Rancona 3.8 FS per 100 pounds of seed.

TREATED SEED LABELING

Seed that has been treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This bag contains seed treated with ipconazole. Do not use for Food, Feed, or Oil Purposes. Persons opening this bag or loading/pouring the treated seed must wear a long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves. After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction."